

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 6:		(11) International Publication Number:	WO 98/33517	
A61K 38/21		(43) International Publication Date:	6 August 1998 (06.08.98)	
 (21) International Application Number: PCT/GB9 (22) International Filing Date: 29 January 1998 (2 (30) Priority Data: 9702021.8 31 January 1997 (31.01.97) (71) Applicant (for all designated States except US): IM COLLEGE OF SCIENCE, TECHNOLOGY & ME [GB/GB]; Sherfield Building, Imperial College, SW7 2AZ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FOSTER, Graha sell [GB/GB]; The Imperial School of Medicin of Medicine, Queen Elizabeth The Queen Mothe St. Mary's Hospital, London W2 1PG (GB). THE Howard, Christopher [GB/GB]; The Imperial School of Medicine, Dept. of Medicine, Queen Elizabeth The Mother Wing, St. Mary's Hospital, London W2 1F (74) Agents: CHAPMAN, Paul, William; Kilburn & Strode Lion Street, London WC1R 4PJ (GB) et al. 	PERIA DICIN Lond am, Ru ee, De er Wir HOMA chool ee Que eG (GI	BY, CA, CH, CN, CU, CZ, DE, GH, GM, GW, HU, ID, IL, IS, LC, LK, LR, LS, LT, LU, LV, MX, NO, NZ, PL, PT, RO, RU, TJ, TM, TR, TT, UA, UG, US, I patent (GH, GM, KE, LS, MW, S patent (AM, AZ, BY, KG, KZ, M patent (AT, BE, CH, DE, DK, E LU, MC, NL, PT, SE), OAPI pa CM, GA, GN, ML, MR, NE, SN Published Published With international search report. Before the expiration of the tin claims and to be republished in amendments.	DK, EE, ES, FI, GB, GE, IP, KE, KG, KP, KR, KZ, MD, MG, MK, MN, MW, SD, SE, SG, SI, SK, SL, UZ, VN, YU, ZW, ARIPO D, SZ, UG, ZW), Eurasian D, RU, TJ, TM), European S, FI, FR, GB, GR, IE, IT, tent (BF, BJ, CF, CG, CI, TD, TG).	

(57) Abstract

The use of an IFN- α subtype is provided for the preparation of a medicament to enhance the T-cell immune response in therapy of cancer, bacterial or parasitic infection or systemic viral infection amongst other disease conditions. Also provided are pharmaceutical formulations which include such sub-types of IFN- α and methods of treatment, including treatment of cancer, bacterial or parasitic infection or systemic viral infection amongst other disease conditions.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia	
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia	
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal	
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland	
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad	
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo	
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan	
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan	
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey	
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago	
BJ	Benin	1E	Ireland	MN	Mongolia	UA	Ukraine	
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda	
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America	
CA	Canada	ıΥ	Italy	MX	Mexico	UZ	Uzbekistan	
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam	
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia	
СН	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe	
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	•		
CM	Cameroon		Republic of Korea	PL	Poland			
CN	China	KR	Republic of Korea	PT	Portugal			
CU	Cuba	KZ	Kazakstan	RO	Romania			
cz	Czech Republic	LC	Saint Lucia	RU	Russian Federation			
DE	Germany	LI	Liechtenstein	SD	Sudan			
DK	Denmark	LK	Sri Lanka	SE	Sweden			
EE	Estonia	LR	Liberia	SG	Singapore	-		

USE OF INTERFERON ALPHA SUBTYPES FOR ENHANCING IMMUNE RESPONSE

The present invention relates to the use of Interferon- α (IFN- α) subtypes, particularly IFN- α_8 , in the preparation of medicaments to treat certain diseases as well as methods of immunomodulation comprising administration of one or more IFN- α subtypes.

Type I interferons (IFN) are a family of closely related glycoproteins containing many IFN- α subtypes and one IFN- β subspecies. At least 13 different human IFN- α subtypes have been identified by analysis of human cDNA libraries and by protein analysis of the IFNs produced by stimulated lymphoblastoid cells. The reasons for this heterogeneity are not yet known. Previous studies have suggested that all Type I IFNs bind to an identical receptor and therefore have identical effects. However a mutant cell line that responds only to IFN- β and interferon- α_8 but not other IFN- α subtypes has been identified showing that these two IFN subspecies either bind to a different receptor or bind in a different way and may therefore have different effects. Molecular analysis of the human Type I IFN receptor thus suggests that the receptor may be able to distinguish between different IFN subtypes.

20

10

A number of studies have compared the effects of different IFN- α subtypes on the antiviral activities of human cell lines. Zoon et al (J. Biol. Chem. 267:15210-

16 (1992) studied IFNs that were obtained from HPLC purification of natural IFN and found no gross differences in their antiviral activities. However, Sperber et al, J. Interferon. Res. 12 363-368 (1992) examined the effects of different recombinant IFN-α subtypes on cells infected with the human immunodeficiency virus (HIV) and found marked differences in their antiviral properties. WO95/24212 disclosed that different IFN-α subtypes were effective antiviral agents in different cell types.

10

Thus, it is possible to target viral infections in say the liver by the use of particular subtypes, eg IFN- α_8 .

B cells or B lymphocytes are a subset of lymphocytes found in secondary lymphoid organs as well as circulating in the blood. They are characterised by the possession of antigen-specific cell surfaceimmunoglobulin molecules of a single antigen-binding specificity which act as receptors for antigen. The interaction of antigen with the cell-surface immunoglobulin is in part responsible for subsequent proliferation of the B cells and their development into antibody-secreting plasma cells. We have found that B cell proliferation can be induced by certain IFN- α subtypes.

T cells are a class of lymphocytes which mediate immune recognition and effect cell-mediated immune responses. In the course of a normal immune response, the binding of a ligand (normally the antigenic complex of peptide and MHC molecule) to the T cell receptor complex (TCR-CD3) on the surface of a T cell initiates intracellular changes, usually leading to proliferation of the T cell concerned and the production of lymphokines.

We have now found that IFN-α subtypes are capable of manipulating the T cell response and particularly enhancing the T cell immune response.

Thus, in a first aspect the present invention provides the use of an IFN- α subtype in the preparation of a medicament to enhance the T cell immune response. In particular, IFN- α_8 is used.

In view of the ability of IFN- α subtypes to enhance the T cell immune response it is possible to use them in the preparation of medicaments to treat certain disese, states. Thus, in further aspects the present invention provides:

- i) the use of an IFN- α subtype in the preparation of a medicament for the treatment of cancer;
- 5 ii) the use of an IFN-α subtype in the preparation of a medicament for the treatment of a bacterial or parasitic infection; and
 - iii) the use of an IFN- α subtype in the preparation of a medicament to treat a systemic viral infection.

Although WO95/24212 disclosed the use of certain IFN- α subtypes in the treatment of viral infections in certain cell types, the present invention relates to their use to treat viral infections which are effectively "systemic", ie viral infections which affect more than one cell or tissue type.

15

Methods for the treatment of the above-noted conditions are also included within the scope of the present invention. Such methods will comprise administration of an effective amount of an IFN- α subtype, in particular IFN- α_8 , to the subject.

- 20 In yet further aspects the present invention provides:
 - i) a pharmaceutical formulation for use in enhancing the T cell immune response which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents;
- 25
- ii) a pharmaceutical formulation for use in the treatment of cancer which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents;

10

20

- iii) a pharmaceutical formulation for use in the treatment of bacterial or parasitic infections which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents; and
- iv) a pharmaceutical formulation for use in the treatment of systemic viral infections which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

In all the above cases the preferred IFN- α subtype is IFN- α_8 .

Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per dose. The precise dose will of course depend on the condition being treated, the route of administration and the age, weight and condition of the patient.

Pharmaceutical compositions within the scope of the present invention may include one or more of the following; preserving agents, solubilising agents, stabilising agents, wetting agents, emulsifiers, sweeteners, colorants, odourants, salts, buffers, coating agents or antioxidants. They may also contain therapeutically active agents.

Pharmaceutical compositions within the scope of the present invention may be adapted for a administration by any appropriate route, for example by the oral (including buccal or sublingual), rectal, nasal, topical (including buccal, sublingual or transdermal), vaginal or parenteral (including subcutaneous, intramuscular, intravenous or intradermal) route. Such a composition may be

prepared by any method known in the art of pharmacy, for example by admixing the active ingredient with a carrier under sterile conditions.

Various routes of administration will now be considered in greater detail:

5

(I) Oral Administration

Pharmaceutical compositions adapted for oral administration may be provided as capsules or tablets; as powders or granules; as solutions, syrups or suspensions (in aqueous or non-aqueous liquids); as edible foams or whips; or as emulsions.

10

Tablets or hard gelatine capsules may comprise lactose, maize starch or derivatives thereof, stearic acid or salts thereof.

Soft gelatine capsules may comprise vegetable oils, waxes, fats, semi-solid, or liquid polyols etc.

Solutions and syrups may comprise water, polyols and sugars. For the preparation of suspensions oils (e.g. vegetable oils) may be used to provide oil-in-water or water-in-oil suspensions.

20

(ii) Transdermal Administration

Pharmaceutical compositions adapted for transdermal administration may be provided as discrete patches intended to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. For example, the active ingredient may be delivered from the patch by iontophoresis (Iontophoresis is described in *Pharmaceutical Research*, 3(6):318 (1986)).

is

25

(iii) Topical Administration

Pharmaceutical compositions adapted for topical administration may be provided

as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols or oils.

For infections of the eye or other external tissues, for example mouth and skin, a topical ointment or cream is preferably used. When formulated in an ointment, the active ingredient may be employed with either a paraffinic or a watermiscible ointment base. Alternatively, the active ingredient may be formulated in a cream with an oil-in-water base or a water-in-oil base.

Pharmaceutical compositions adapted for topical administration to the eye include eye drops. Here the active ingredient can be dissolved or suspended in a suitable 10 carrier, e.g. in an aqueous solvent.

Pharmaceutical compositions adapted for topical administration in the mouth include lozenges, pastilles and mouthwashes.

15 .

5

(iv) Rectal Administration

Pharmaceutical compositions adapted for rectal administration may be provided as suppositories or enemas.

20 (v) Nasal Administration

Pharmaceutical compositions adapted for nasal administration which use solid carriers include a coarse powder (e.g. having a particle size in the range of 20 to 500 microns). This can be administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nose from a container of powder held close to the nose.

25

Compositions adopted for nasal administration which use liquid carriers include nasal sprays or nasal drops. These may comprise aqueous or oil solutions of the active ingredient.

Pharmaceutical compositions adapted for administration by inhalation include fine particle dusts or mists, which may be generated by means of various types of apparatus, e.g. pressurised aerosols, nebulizers or insufflators. Such apparatus can be constructed so as to provide predetermined dosages of the active ingredient.

(vi) Vaginal Administration

Pharmaceutical compositions adapted for vaginal administration may be provided as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

(vii) Parenteral Administration

Pharmaceutical compositions adapted for parenteral administrations include aqueous and non-aqueous sterile injectable solutions or suspensions. These may contain antioxidants, buffers, bacteriostats and solutes which render the compositions substantially isotonic with the blood of an intended recipient. Other components which may be present in such compositions include water, alcohols, polyols, glycerine and vegetable oils, for example. Compositions adapted for parenteral administration may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of a sterile liquid carrier, e.g. sterile water form injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets.

25

20

Preferred unit dosage formulations are those containing a daily dose or sub-dose, as herein above recited, or an appropriate fraction thereof, of an active ingredient.

In a final aspect the present invention provides a method for enhancing the T cell

immune response which comprises administering to a subject an effective amount of an IFN- α subtype.

The invention will now be described by reference to the following example which should not be construed as in any way limiting the scope of the invention.

The examples refer to the figures in which:

FIGURE 1: shows inhibition of peripheral blood T cell proliferation in response to IL-2 and anti-CD3; and

FIGURE 2: shows the effect of various IFN α subtypes on the production of IFN γ by T cells in the presence or absence of anti-CD3 antibodies plus IL-2.

15 .

20

25

EXAMPLE 1

Peripheral blood T cells were purified by Ficoll density gradient centrifugation and E rosetting with sheep red blood cells (SRBC) to separate B and T cells. The T cells were recovered from the rosettes by lysis of the SRBC and were cultured at 1x106 cells/ml in RPMI 1640 medium with 10% FCS and gentamycinfor three days.

³H thymidine was added for the last 8 hours culture and incorporation was measured by scintilllation counting. The cells were stimulated with anti-CD3 (UCHT1) at 2μg/ml. Under these conditions none of the IFN-α subtypes induced significant proliferation of the cells (data not shown).

However, when the T cells were stimulated with anti-CD3 and IL-2, inhibition of the IL-2 induced proliferation could be seen (see figure 1). IFN- α_8 was the most

effective, with the other subtypes inhibiting to a lesser extent with the exception of IFN- α^1 which was inactive in the assay.

EXAMPLE 2

Peripheral blood T cells were purified as in example 1. To investigate stimulation the cells were exposed to IFNα subtypes in the presence or absence of anti-CD3 antibodies (UCHT-1) plus IL-2 and IFNγ production was measured using standard intracellular staining techniques. The results are shown in figure 2 and indicate that anti-CD3 plus IL-2 antibodies plus IFNα₈ caused an increase in the proportion of cells producing IFNγ.

WO 98/33517 PCT/GB98/00269

10

CLAIMS:

1. The use of an IFN- α subtype in the preparation of a medicament to enhance the T cell immune response.

5


- 2. The use of an IFN- α subtype in the preparation of a medicament for the treatment of cancer.
- 3. The use of an IFN- α subtype in the preparation of a medicament for the treatment of a bacterial or parasitic infection.
 - 4. The use of an IFN- α subtype in the preparation of a medicament to treat a systemic viral infection.
- 15 5. The use as claimed in any one of claims 1 to 4 wherein the subtype is IFN- α_8 .
 - 6. A pharmaceutical formulation for use in enhancing the T cell immune response which comprises an IFN-α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

20


- 7. A pharmaceutical formulation for use in the treatment of cancer which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.
- 8. A pharmaceutical formulation for use in the treatment of bacterial or parasitic infections which comprises an IFN-α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.
 - 9. A pharmaceutical formulation for use in the treatment of systemic viral

infections which comprises an IFN- α subtype, optionally together with one or more pharmaceutically acceptable carriers, excipients or diluents.

- 10. A pharmaceutical formulation as claimed in any one of claims 6 to 9 wherein 5 the subtype is IFN- α_8 .
 - 11. A method of enhancing a subject's T cell immune response which comprises administering an effective amount of an IFN- α subtype to the subject.
- 10 12. A method for the treatment of cancer which comprises administering an effective amount of an IFN- α subtype to the subject.
 - 13. A method for the treatment of a bacterial or parasitic infection which comprises administering an effective amount of an IFN- α subtype to the subject.
 - 14. A method for the treatment of a systemic viral infection which comprises administering an effective amount of an IFN- α subtype to the subject.
- 15. A method as claimed in any one of claims 11 to 14 wherein the subtype is 20 IFN- α_8 .



SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)

Internat : Application No PCT/GB 98/00269

IPC 6	FICATION OF SUBJECT MATTER A61K38/21		
According to	nternational Patent Classification (IPC) or to both national classifica	tion and IPC	
	SEARCHED		
IPC 6	noumentation searched (classification system followed by classification A61K	n symbols)	
Documentat	tion searched other than minimum documentation to the extent that su	ich documents are included in the fields search	hed
Electronic d	ata base consulted during the international search (name of data bas	e and, where practical, search terms used)	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the rele	vant passages	Relevant to claim No.
. Y	WO 95 24212 A (IMPERIAL COLLEGE OF SCIENCE, TECHNOLOGY & MEDICINE) IS September 1995 cited in the application see the whole document		1-15
)y •	WO 94 14474 A (SCHERING CORPORAT) July 1994 see the whole document	1-15	
A	WO 94 20122 A (GEORGETOWN UNIVERS September 1994 see the whole document	SITY) 15	1-15
		-/	
X Furt	her documents are listed in the continuation of box C.	X Patent family members are listed in a	unnex.
"A" docume consid "E" earlier filing c	stegories of cited documents : ent defining the general state of the art which is not dered to be of particular relevance document but published on or after the international date in the document but published on priority olaim(s) or is cited to establish the publication date of another	"T" later document published after the interns or priority date and not in conflict with the cited to understand the principle or theor invention. "X" document of particular relevance; the claim cannot be considered novel or cannot be involve an inventive step when the document of particular relevance; the claim to document of particular relevance; the claim to the considered novel or the document of particular relevance; the claim to the considered novel or	a application but y underlying the med invention a considered to ment is taken alone
oitatio "O" docum other "P" docum	n or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or means ent published prior to the international filling date but han the priority date claimed	cannot be considered to involve an inver- document is combined with one or more ments, such combination being obvious in the art. *& document member of the same patent far	ntive step when the other such docu- to a person skilled
	actual completion of the international search	Date of mailing of the international search 2 9. 05.	
	1 May 1998	20.00	1000
Name and	mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nt, Fax: (+31-70) 340-3016	Authorized officer Moreau, J	

1

Internativ Application No PCT/GB 98/00269

	WALL DOGGET TO DE DEL EVANT	PC1/GB 98	700203	
C.(Continua Category *	tion) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No.	
Category			<u> </u>	
A	FOSTER G R ET AL: "Different relative activities of human cell-derived interferon -alpha subtypes: IFN-alpha has very high antiviral potency." JOURNAL OF INTERFERON AND CYTOKINE RESEARCH 16 (12). 1996. 1027-1033, XP002064516 see the whole document		1-15	
P,X	HIBBERT L M ET AL: "Activity of different interferon alpha subtypes: alpha-8 is the most potent anti-viral subtype and has unique immunomodulatory properties." 32ND ANNUAL MEETING OF THE EUROPEAN ASSOCIATION FOR THE STUDY OF LIVER, LONDON, ENGLAND, UK, APRIL 9-12, 1997. JOURNAL OF HEPATOLOGY 26 (SUPPL. 1). 1997. 186, XP002064517 see the whole document		1-15	
	See the whole document			
•				
	·			
			-	
		•		

Inten Junal application No. PCT/GB 98/00269

Boxi	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This Inte	ernational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
	Although claims 11-15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
з. 🗌	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This Int	emational Searching Authority found multiple inventions in this international application, as follows:
	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	k on Protest The additional search fees were accompanied by the applicant's protest.
	No protest accompanied the payment of additional search fees.

Into mation on patent family members

Internati Application No PCT/GB 98/00269

Patent document cited in search repor	1	Publication date		atent family nember(s)	Publication date
WO 9524212	A	14-09-1995	AU EP JP	1854695 A 0741577 A 9509955 T	25-09-1995 13-11-1996 07-10-1997
WO 9414474	Α	07-07-1994	AU MX	5953494 A 9307733 A	19-07-1994 30-06-1994
WO 9420122	Α	15-09-1994	AU	6354994 A	26-09-1994

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
□ OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.